

Article



Gillies, George

TN#: 1818299



User ID:



Article

Article

Call #: QC1 .A56 ser.2 v.19

Jan-June 1974

X002409317

Preferred Delivery: **Library**

Dept: **Engineering**

Address: **Mech. Eng.**

**105D**

Pick up Library: **PHYS**

Location: **IVY BY-REQUEST**

Book/Journal Title:

**BULLETIN OF THE AMERICAN PHYSICAL SOCIETY**

Book Author:

Other Info:

Volume: **19**

Issue: **5**

Year: **1974**

Pages: **688-689**

Article Author: **BEAMS, JW**

Article Title: **MAGNETIC SUSPENSION OSMOMETER**

Document Delivery - Ivy Stacks/LEO - Article

Status: **Faculty**

Email Address: **gtg@virginia.edu**

VIRGO:

Copyright Information:

NOTICE: This material may be protected by copyright law (Title 17, United States Code)

Policy:

11/27/2018 12:31:51 PM (cdh2n) Item ID: X002409317

Date Needed: 02/25/2019

Email Address: **gtg@virginia.edu**

George Gillies

Pick up Library: **PHYS**

University of Virginia  
Alderman Library  
Interlibrary Services  
PO Box 400109  
160 N. McCormick Road  
Charlottesville, VA 22904-4109

(EDWARD F. ZGANJAR, Louisiana State University, presiding)

## General Interest

INVITED PAPER (30 minutes)

CB1. Are there Viable Alternatives to Nuclear Power? P.F. ZWEIFEL and ANITA GUYNN, Virginia Polytechnic Institute and State U.

INVITED PAPER (30 minutes)

CB2. Prospects for Physics in Energy Research. ALEX ZUCKER, Oak Ridge National Laboratory.

INVITED PAPER (30 minutes)

CB3. On Some of Physic's Problems. EUGENE P. WIGNER, Princeton University. -- Two kinds of problems will be discussed: the problems of the general progress of physics and its relations with society; and some true scientific problems. A very serious problem of the first kind is the difficulty the physicist experiences when he wishes to maintain contact with the whole body

of his discipline and to avoid excessive specialisation. The second problem is actually the result of the success of physics, the natural sciences, and technology. This success makes it physically possible, at least in countries where it is fully operative, for everyone to have a materially carefree life. Hence, new developments in physics and technology are less needed; society's problems became of a social and human nature.

Of scientific problems only two will be discussed; these two suggest that present day physics is not an all-compassing science. The first problem concerns the age of the universe, why Wammetod has not arrived yet. The implications of the standard, statistical interpretation of quantum mechanics present the second problem. In this connection, some attention will be given to certain technical weaknesses of this interpretation.

FRIDAY MORNING AT 8:30 SESSION CC PARLOR D

(PERRY MORTON, Samford University, presiding)

## Biological Physics

C C1. Energy Levels of Transition Metal Ions in Biologically Active Sites; Arthur S. Brill, University of Virginia (30 min.)

INVITED PAPER (30 minutes)

CC2. Free Radical Reactions in Irradiated Organic Single Crystals. G. C. MOULTON, Florida State University. -- The free radicals produced by irradiation in single crystals of several amino acids, dicarboxylic acids, and amides using electron spin resonance and electron nuclear double resonance spectroscopy for detection and analysis will be discussed. Some free radical reaction mechanisms and pathways will be presented. It has been found that the particular pathway that a free radical reaction follows can in some cases be sensitive to a number of parameters including the crystal form and also whether the crystal was irradiated at low temperature and warmed to room temperature or irradiated at room temperature.

CC3. Creation and Destruction of a Radical in Guanine by X-rays. D. L. ALLISON, Florence State University, and C. ALEXANDER, JR., The University of Alabama.

CC4. A Comparison of the Energy Requirements and Broad Spectral Sensitivity of the Melanophores of the Fiddler Crab. T. P. COOHILL, Western Kentucky University, Bowling Green, Kentucky -- The broad spectral sensitivity to light of one of the color cells of the fiddler crab, Uca pugilator, was investigated. The primary response (direct effect) of these cells was most sensitive to near ultraviolet radiation (3000 + 4000Å). The secondary response (hormonal) was sensitive to all spectral regions tested

(2000 + 7000Å). Less energy is required to elicit the secondary response in these crabs. Even the output of a microscope lamp was sufficient to invoke a strong secondary response. The adaptive value of this will be discussed.

\*Supported in part by a Faculty Research Grant from WKU and by NSF grant #GB-27497.

† Submitted by WILLIAM G. BUCKMAN

CC5. Magnetic Suspension Osmometer.\* J.W. BEAMS, M.G. HODGINS, and D.W.KUPKE, Univ. of Virginia -- Very small osmotic pressures are measured by a modified magnetic suspension balance method previously used for measuring densities and viscosities of protein and other solutions.<sup>1</sup> A small hollow ferromagnetic cylinder (buoy) closed at the top and open at the bottom and filled with trapped air is freely magnetically suspended below the surface of the solution. The solution is contained in a 0.2 CC completely filled glass tube coaxial with the vertical axis of the buoy and closed at the bottom with a semi-permeable membrane and at the top with a liquid tight valve which may be opened or closed. This glass container is completely immersed in the solvent. Because of the trapped air, the buoy functions like a Cartesian diver and is pressure sensitive. The method is especially useful for determining osmotic pressures in small samples of dilute

solutions ( $\sim 10^{-6}M$ ) or of solutions of macromolecules where the pressure varies from 1 cm to  $\sim 10^{-4}$  cm of water.

\*Supported by USPH-NIGM Grant GM19682-01 and NSF Grants GP-31721, GB-27331.  
D.W. Kupke, M.G. Hodgins, and J. W. Beams, Proc. Natl. Acad. Sci. USA 69, 2258 (1972).

CC6. Magnetic Suspension Densimeter-Viscometer.\* M.G. HODGINS, D. W. KUPKE, and J. W. BEAMS, Univ. of Virginia -- The magnetic suspension densimeter-viscometer previously described<sup>1</sup> is here modified so that the simultaneous measurement of the density and viscosity of a solution may be made  $\sim 10^2$  times per sec. A small cylindrical buoy is magnetically suspended inside of and on the axis of a sealed glass cylinder which is completely filled with the solution under investigation. The vertical glass cylinder is slowly rotated around its vertical axis by an accurate clock mechanism and the buoy is held stationary by a rotating magnetic field. The power required to hold

the buoy stationary gives the viscosity, since the magnetic suspension itself exerts no restoring axial torque on the buoy, while the current in the solenoids which support the buoy is a measure of the density. The measurements are made on samples of solutions of 0.2 CC. The use of the apparatus to follow physical changes, chemical reactions, etc. and the precision obtained will be discussed.

\*Supported by NSF Grants GP-31721, GB-27331 and USPH-NIGM Grant GM19682-01.

<sup>1</sup>D.W. Kupke, M.G. Hodgins, and J.W. Beams, Proc. Natl. Acad. Sci. USA 69, 2258 (1972).

CC7. A Preparation and Initiation of a Comparative Study of Hemolysate Prepared from Normal and Sick Cell Hemoglobin Using Mossbauer Spectroscopy.\* Bruce L. Biskey and Donald E. Kinkaid, North Georgia College

FRIDAY MORNING AT 8:30 SESSION CD PARLOR E

(BARLOW NEWBOLT, Washington and Lee University, presiding)

## Solid State Physics II

CD1. Computer Simulation of the Structural Deformation of an Ice Surface. DONALD E. HAGEN and JAMES L. KASSNER, JR., Univ. of Missouri-Rolla -- We use a surface energy minimization technique to develop a model for the structure of an ice surface. We consider the special case of an ice crystal that is cleaved perpendicular to the c-axis. The resulting singly-bonded surface molecules are allowed to move so they can hydrogen bond with one another and thereby lower the surface energy. This movement is opposed by the strain that arises when either the bonds to the lower molecular layers or the newly formed bonds are stretched or bent. Movement can occur because the hydrogen bond energy is great enough to support substantial structural strain energy before dissociation. The structure is allowed to relax until the sum of the negative binding energy and positive strain energy reaches a minimum. It is found that substantial deviation from the perfect ice lattice exists at the surface.

CD2 Thermal Expansion Measurements in a 500-degree Temperature Range. J. S. BROWDER\* and S. S. BALLARD, Univ. of Florida

CD3 Interstitial Migration in Mo. R. R. COLTMAN, JR., C. E. KLABUNDE, J. K. REDMAN, and A. L. SOUTHERN, Oak Ridge National Laboratory.

CD4. A Mössbauer Study of Disordered Iron-Nickel-Aluminum Alloys.\* M. E. SANDERS and B. DE MAYO, West Georgia Coll. -- Room temperature, <sup>57</sup>Fe Mössbauer spectra have been obtained for atomically disordered fcc (Fe<sub>x</sub>Ni<sub>1-x</sub>)<sub>y</sub>Al<sub>1-y</sub> with x = 0.17 and y = 0.85, 0.90, 0.95 and 1.0; and with y = 0.90 and x = 0.05, 0.17, 0.33, and 0.45. Values were obtained for the magne-

tic hyperfine field at the iron nuclei and for the iron isomer shifts, quadrupole splittings and line shapes. The hyperfine field varies from 282 + 2 kOe for Fe<sub>0.17</sub>Ni<sub>0.83</sub> to 183 + 5 kOe for (Fe<sub>0.17</sub>Ni<sub>0.83</sub>)<sub>0.85</sub>Al<sub>0.15</sub>. The isomer shifts and quadrupole splittings are small and comparable to those of similar alloys. The line shapes indicated that aluminum has a shorter range effect on the iron atoms than does nickel. The rigid band model fails to adequately explain the results.

\*Work supported in part by the Faculty Research Committee of West Georgia College.

CD5. General Solutions of Radiation Defect Annealing Equations for Elemental Semiconductors. W.M. Pritchard, Old Dominion U. and J.L.Reid, Clemson U. -- General solutions are presented in power series form of a set of kinetic equations for the isothermal annealing of simple-radiation defects in elemental semiconductors. These equations describe the case where the initial damage state is associated with uncorrelated vacancies and interstitials and the annealing process involves trapping and annihilation of the mobile defect. A kinetic model with an arbitrary number of types of secondary defect complexes is considered as well as a simple model<sup>1</sup> based on only one species of defect complex. The series solutions are well suited for computer evaluation of numerical results. For the simple annealing model with parameters characteristic of silicon and germanium, truncation errors are easily estimated since the terms have alternating signs. For cases where the total annealing time is greater than the intervals of series convergence, a repetitive series expansion approach can be employed.

<sup>1</sup>A.C. Damask and G.J. Dienes, Phys. Rev. 125,444(1962).



(EDWARD F. ZGANJAR, Louisiana State University, presiding)

## General Interest

INVITED PAPER (30 minutes)

CB1. Are there Viable Alternatives to Nuclear Power? P. F. ZWEIFEL and ANITA GYNN, Virginia Polytechnic Institute and State U.

INVITED PAPER (30 minutes)

CB2. Prospects for Physics in Energy Research. ALEX ZUCKER, Oak Ridge National Laboratory.

INVITED PAPER (30 minutes)

CB3. On Some of Physic's Problems. EUGENE P. WIGNER, Princeton University. -- Two kinds of problems will be discussed: the problems of the general progress of physics and its relations with society; and some true scientific problems. A very serious problem of the first kind is the difficulty the physicist experiences when he wishes to maintain contact with the whole body

of his discipline and to avoid excessive specialisation. The second problem is actually the result of the success of physics, the natural sciences, and technology. This success makes it physically possible, at least in countries where it is fully operative, for everyone to have a materially carefree life. Hence, new developments in physics and technology are less needed; society's problems became of a social and human nature.

Of scientific problems only two will be discussed; these two suggest that present day physics is not an all-compassing science. The first problem concerns the age of the universe, why Warmedot has not arrived yet. The implications of the standard, statistical interpretation of quantum mechanics present the second problem. In this connection, some attention will be given to certain technical weaknesses of this interpretation.

(PERRY MORTON, Samford University, presiding)

## Biological Physics

C C1. Energy Levels of Transition Metal Ions in Biologically Active Sites; Arthur S. Brill, University of Virginia (30 min.)

INVITED PAPER (30 minutes)

CC2. Free Radical Reactions in Irradiated Organic Single Crystals. G. C. MOULTON, Florida State University. -- The free radicals produced by irradiation in single crystals of several amino acids, dicarboxylic acids, and amides using electron spin resonance and electron nuclear double resonance spectroscopy for detection and analysis will be discussed. Some free radical reaction mechanisms and pathways will be presented. It has been found that the particular pathway that a free radical reaction follows can in some cases be sensitive to a number of parameters including the crystal form and also whether the crystal was irradiated at low temperature and warmed to room temperature or irradiated at room temperature.

CC3. Creation and Destruction of a Radical in Guanine by X-rays. D. L. ALLISON, Florence State University, and C. ALEXANDER, JR., The University of Alabama.

CC4. A Comparison of the Energy Requirements and Broad Spectral Sensitivity of the Melanophores of the Fiddler Crab. T. P. COOHILL, Western Kentucky University, Bowling Green, Kentucky -- The broad spectral sensitivity to light of one of the color cells of the fiddler crab, Uca pugnator, was investigated. The primary response (direct effect) of these cells was most sensitive to near ultraviolet radiation (3000 + 4000Å). The secondary response (hormonal) was sensitive to all spectral regions tested

(2000 + 7000Å). Less energy is required to elicit the secondary response in these crabs. Even the output of a microscope lamp was sufficient to invoke a strong secondary response. The adaptive value of this will be discussed.

\*Supported in part by a Faculty Research Grant from WKU and by NSF grant #GB-27497.

† Submitted by WILLIAM G. BUCKMAN

CC5. Magnetic Suspension Osmometer.\* J.W. BEAMS, M.G. HODGINS, and D.W.KUPKE, Univ. of Virginia -- Very small osmotic pressures are measured by a modified magnetic suspension balance method previously used for measuring densities and viscosities of protein and other solutions.<sup>1</sup> A small hollow ferromagnetic cylinder (buoy) closed at the top and open at the bottom and filled with trapped air is freely magnetically suspended below the surface of the solution. The solution is contained in a 0.2 CC completely filled glass tube coaxial with the vertical axis of the buoy and closed at the bottom with a semi-permeable membrane and at the top with a liquid tight valve which may be opened or closed. This glass container is completely immersed in the solvent. Because of the trapped air, the buoy functions like a Cartesian diver and is pressure sensitive. The method is especially useful for determining osmotic pressures in small samples of dilute

solutions ( $\sim 10^{-6}M$ ) or of solutions of macromolecules where the pressure varies from 1 cm to  $\sim 10^{-4}$  cm of water.

\*Supported by USPH-NIGM Grant GM19682-01 and NSF Grants GP-31721, GB-27331.

1p.W. Kupke, M.G. Hodgins, and J. W. Beams, Proc. Natl. Acad. Sci. USA 69, 2258 (1972).

CC6. Magnetic Suspension Densimeter-Viscometer.\* M.G. HODGINS, D. W. KUPKE, and J. W. BEAMS, Univ. of Virginia -- The magnetic suspension densimeter-viscometer previously described<sup>1</sup> is here modified so that the simultaneous measurement of the density and viscosity of a solution may be made  $\sim 10^2$  times per sec. A small cylindrical buoy is magnetically suspended inside of and on the axis of a sealed glass cylinder which is completely filled with the solution under investigation. The vertical glass cylinder is slowly rotated around its vertical axis by an accurate clock mechanism and the buoy is held stationary by a rotating magnetic field. The power required to hold

the buoy stationary gives the viscosity, since the magnetic suspension itself exerts no restoring axial torque on the buoy, while the current in the solenoids which support the buoy is a measure of the density. The measurements are made on samples of solutions of 0.2 CC. The use of the apparatus to follow physical changes, chemical reactions, etc. and the precision obtained will be discussed.

\*Supported by NSF Grants GP-31721, GB-27331 and USPH-NIGM Grant GM19682-01.

1p.W. Kupke, M.G. Hodgins, and J.W. Beams, Proc. Natl. Acad. Sci. USA 69, 2258 (1972).

CC7. A Preparation and Initiation of a Comparative Study of Hemolysate Prepared from Normal and Sick Cell Hemoglobin Using Mossbauer Spectroscopy.\* Bruce L. Biskey and Donald E. Kinkaid, North Georgia College

(BARLOW NEWBOLT, Washington and Lee University, presiding)

## Solid State Physics II

CD1. Computer Simulation of the Structural Deformation of an Ice Surface. DONALD E. HAGEN and JAMES L. KASSNER, JR., Univ. of Missouri-Rolla -- We use a surface energy minimization technique to develop a model for the structure of an ice surface. We consider the special case of an ice crystal that is cleaved perpendicular to the c-axis. The resulting singly-bonded surface molecules are allowed to move so they can hydrogen bond with one another and thereby lower the surface energy. This movement is opposed by the strain that arises when either the bonds to the lower molecular layers or the newly formed bonds are stretched or bent. Movement can occur because the hydrogen bond energy is great enough to support substantial structural strain energy before dissociation. The structure is allowed to relax until the sum of the negative binding energy and positive strain energy reaches a minimum. It is found that substantial deviation from the perfect ice lattice exists at the surface.

CD2. Thermal Expansion Measurements in a 500-degree Temperature Range. J. S. BROWDER\* and S. S. BALLARD, Univ. of Florida

CD3. Interstitial Migration in Mo. R. R. COLTMAN, JR., C. E. KLABUNDE, J. K. REDMAN, and A. L. SOUTHERN, Oak Ridge National Laboratory.

CD4. A Mössbauer Study of Disordered Iron-Nickel-Aluminum Alloys.\* M. E. SANDERS and B. DE MAYO, West Georgia Coll. -- Room temperature, <sup>57</sup>Fe Mössbauer spectra have been obtained for atomically disordered fcc (Fe<sub>x</sub>Ni<sub>1-x</sub>)<sub>2</sub>Al<sub>1-y</sub> with x = 0.17 and y = 0.85, 0.90, 0.95 and 1.0; and with y = 0.90 and x = 0.05, 0.17, 0.33, and 0.45. Values were obtained for the magne-

tic hyperfine field at the iron nuclei and for the iron isomer shifts, quadrupole splittings and line shapes. The hyperfine field varies from 282 + 2 kOe for Fe<sub>0.17</sub>Ni<sub>0.83</sub> to 183 + 5 kOe for (Fe<sub>0.17</sub>Ni<sub>0.83</sub>)<sub>0.85</sub>Al<sub>0.15</sub>. The isomer shifts and quadrupole splittings are small and comparable to those of similar alloys. The line shapes indicated that aluminum has a shorter range effect on the iron atoms than does nickel. The rigid band model fails to adequately explain the results.

\*Work supported in part by the Faculty Research Committee of West Georgia College.

CD5. General Solutions of Radiation Defect Annealing Equations for Elemental Semiconductors. W.M. Pritchard, Old Dominion U. and J.L.Reid, Clemson U. -- General solutions are presented in power series form of a set of kinetic equations for the isothermal annealing of simple-radiation defects in elemental semiconductors. These equations describe the case where the initial damage state is associated with uncorrelated vacancies and interstitials and the annealing process involves trapping and annihilation of the mobile defect. A kinetic model with an arbitrary number of types of secondary defect complexes is considered as well as a simple model<sup>1</sup> based on only one species of defect complex. The series solutions are well suited for computer evaluation of numerical results. For the simple annealing model with parameters characteristic of silicon and germanium, truncation errors are easily estimated since the terms have alternating signs. For cases where the total annealing time is greater than the intervals of series convergence, a repetitive series expansion approach can be employed.

<sup>1</sup>A.C. Damask and G.J. Dienes, Phys. Rev. 125,444(1962).



MAY 17 1974

# bulletin


OF THE AMERICAN PHYSICAL SOCIETY

MAY 1974

INCLUDING THE PROGRAM OF THE  
1974 JUNE MEETING IN SALT LAKE CITY, UTAH  
12-14 JUNE 1974

PUBLISHED FOR THE AMERICAN PHYSICAL SOCIETY  
BY THE AMERICAN INSTITUTE OF PHYSICS

NVI815220 BAPS 019005 10  
VIRGINIA, UNIV OF CHARLOTTE  
VILLE ALDERMAN LIB-SERS-PERIOD  
CHARLOTTESVILLE VA 22901



The Sheraton Park Hotel

Phys  
CC  
1  
456  
472  
413